

trees will probably need more fertilizer this year applied more often. Young trees should receive small amounts of fertilizer applied monthly. University trials with phosphorus and potassium did not indicate any consistent trends for yield increase; the advice here is to apply these fertilizers only if the trees show deficiencies in the leaf analysis.

Weed Control

Spot treatments with Roundup should control most of the problems. If there are large areas of soil exposed to the sun, an application of a weed seed sterilant such as Princep Caliber 90 should be applied now.

Tree Thinning or Block Stumping

If the grove is crowded, now is the time for this good cultural practice. Trees should be stumped at about 5 feet in height; this allows you to have a man with a chain saw standing on the ground rather than on a ladder. Remember to harvest the trees before cutting and paint the trunks with a water-based white paint for sunburn protection.

Tree Planting

March is the best time to plant replacement trees. Try to avoid replanting in root-rot infested areas; it would be better to replant with citrus. If you wish to replant avocados, make sure the rootstock is Duke 7, Thomas or Toro Canyon and the sleeves have been drenched with Aliette or Ridomil before planting.

**Only
California
Grower
Covers The
Avocado Industry.
Stay Informed.
To Subscribe Call
1 — 800
678 — GROW**

Does An Avocado Early Bloom Spray Make Dollars And Sense?

by Carol J. Lovatt

The avocado is a seeded fruit. Therefore, pollination and fertilization are both critical to fruit set and final yield. Pollination is the arrival of the pollen (the male part of the reproductive cycle) on the stigma of the pistil

(the female part of the flower). Under optimal conditions, the pollen germinates producing a pollen tube which grows through the tissues of the stigma, style, and ovary to the ovule,

which contains the egg. The pollen tube delivers the sperm to the egg. The fusion of sperm and egg is fertilization. The product of fertilization is the embryo, which develops into the young avocado seedling within the ovule. After fertilization, the ovule develops into the seed within the ovary, which develops into the avocado fruit. The successful completion of these two processes is essential to fruit set.

The goal of our research is to increase fruit set and yield by improving the probability that fertilization will occur. Our approach is to lengthen the duration of the effective pollination

period by increasing the growth rate of the pollen tubes and extending the period of ovule viability. The finite period of time during which fertilization can successfully take place is called the effective pollination period. Conceptually and mathematically, the effective pollination period of a crop is the longevity of the ovule minus the length of time required for the pollen tube to reach the ovule to deliver the sperm to the egg so that fertilization can take place.

Temperatures prevailing during bloom are known to influence the effective pollination period of a seeded-fruit crop, such as avocado, and thus influence fruit set and yield. Cool temperatures which prevail during the flowering



A single application of boron to the canopy during early bloom, pictured here, resulted in 126 more pounds of fruit per tree.

period decrease the viability of the ovule and increase the length of time it takes for the pollen tubes to grow from the stigma to the ovule. Thus, the duration of the effective pollination period is significantly shortened and fruit set is reduced. Warm temperatures during flowering increase both ovule longevity and the growth rate of the pollen tubes. This correspondingly increases the effective pollination period and thereby fruit set.

Ovule viability has been improved in deciduous tree crops by a summer application of nitrogen despite the fact that the trees were not nitrogen-deficient. The use of nitrogen to enhance ovule viability, fruit set and yield in avocado has not been investigated.

Carol J. Lovatt is an Associate Professor of Plant Physiology at University of California, Riverside.

For more information circle #4 on reader service card

FRUIT GROWERS SUPPLY COMPANY

Whatever the crop, whatever the need, stop in or call FGS for supplies and service

FGS SERVICES:

Full stock of agricultural crop supplies
On-site buyers for special order services
Irrigation systems & material sales

ALL PRODUCTS AVAILABLE AT COMPETITIVE PRICES.

Agricultural Management Supplies
Agricultural Chemicals
Fertilizer and Fertilizer Handling Equipment
Irrigation System Design, Installation & Material
Machinery and Equipment
Pruning Materials
Safety Equipment
Weather and Temperature Recording Instruments
Harvesting Equipment and Supplies

FRUIT GROWERS SUPPLY COMPANY



Orange Cove	(209) 626-4629
Porterville	(800) 354-2355
Woodlake	(800) 637-1003
Riverside	(800) 323-8303
Santa Paula	(800) 528-9347
Yuma, AZ	(602) 726-0250

avocados

It has been known for a long time that boron is essential for pollen germination, for successful growth of a pollen tube through the stigma, style, and ovary to the ovule, and for the mitotic divisions necessary to produce the sperm. Boron applied during fall or spring to trees that were not boron deficient increased fruit set in a number of deciduous fruit tree crops, especially when cooler temperatures prevailed during bloom. An investigation of the effect of boron on fruit set of the 'Hass' avocado has recently been initiated in South Africa. Fruit set and yield were improved, although not significantly, by spraying trees with 0.27 oz. Solubor per gallon of water at the early stages of flowering.

On the basis of these findings, we designed a field trial to test the hypothesis that fruit set and yield of avocado can be improved by increasing the effective pollination period by prolonging ovule viability with a bloom application of nitrogen and accelerating pollen tube growth rate with a bloom application of boron.

Warren Currier, Secretary of the California Avocado Development Organization, was the driving force behind this research, which was funded for the first two years by CADO. The research was initiated in collaboration with Dr. Howard Ohr, Cooperative Extension Plant Pathologist, and Dr. Guy Witney, Farm Advisor, University of California, Riverside. We have completed two years of field research, will harvest the third year this June, and will continue the research for two additional years with matching funds from the California Avocado Society and the California Department of Food and Agriculture Fertilizer Research and Education Program. We are indebted to Dick Whiley of AgriSpec and Chris Taylor of the Limoneira Company for their cooperation and use of the Hass avocado orchards.

The first year of the field experiment was a screening trial employing only 8 individual tree replicates per treatment. The optimum treatment conditions

determined in the first year were employed in the second and subsequent years in well-replicated experiments employing 16 individual tree replicates per treatment.

Nitrogen was applied to the canopy as ~~16~~ ^{1.6 pini} pint Unocal Plus, liquid low-biuret urea (~~20~~ ¹⁰ percent N, 0.1 percent biuret) in 4 gallons of water at pH 5.5 to 6.5. This application rate supplies 6 oz. (0.375 lbs.) N to the tree.

Boron was applied to the canopy as 1.06 oz. (0.066 lbs.) Solubor, granular sodiumtetraborate (20.5 percent B) in 4 gallons of water at pH 5.5 to 6.5. This application rate supplies 0.22 oz. (0.014 lbs) B to the tree.

Maximum increase in yield was obtained by spraying the canopy when approximately 50 percent of the trees in the orchard had 50 percent of their inflorescences at chest height at the early bloom stage of inflorescence development, approximately mid-March. Inflorescences on some trees, especially at the top of the tree, were more advanced. A positive but less dramatic response in yield was obtained when the majority of inflorescences were at a more advanced stage, approximately mid-April. Application of nitrogen or boron at budbreak (early January) did not have any affect on yield.

Although not significant at the 5 percent level, for the first year of the study (an "off" year) the single application of low-biuret urea to the canopy during early bloom resulted in 75 lbs. more fruit per tree, a 50 percent increase in yield over the untreated control trees. This resulted in an increase of ~~75~~ ⁷⁶ per tree net return to the grower. Two canopy sprays of boron, one at early bloom and a second approximately 30 days later when the inflorescences were more expanded, were required to increase yield over the control trees, but the increase was only half that obtained with the single early bloom spray of urea.

In the second year of the study (an "on" year), the single application of boron to the canopy during early bloom resulted in 126 lbs. more fruit per tree, a

For more information circle #36 on reader service card

ORDER NOW!

CITRUS
MAJOR VARIETIES
LEADING ROOTSTOCKS
AVOCADOS
CLONAL AND SEEDLING
ROOTSTOCKS



WE SHIP TO ALL CALIF. COUNTIES

C & M NURSERY

277 W. TEFFT • P.O. BOX 383
NIPOMO, CALIFORNIA 93444
PHONE 805-929-1941
FAX 805-929-5588

50 percent increase in yield over the untreated control trees ($P < 0.05$). This resulted in an increase of \$40 per tree net return to the grower. No other treatment significantly increased yield over the control.

Two additional results obtained from this research must be noted. First, there was a significant negative interaction resulting in a reduction in yield for trees receiving early bloom sprays of both urea and boron, regardless of whether urea was applied simultaneously with boron or 72 hours prior to boron. Second, use of trunk injections of urea or boron to raise the tree nutrient status at early bloom without spraying the developing flowers provided strong evidence that the nutrients must be applied directly onto the developing flowers. Trunk injections successfully increased the nutrient content of the leaves but did not increase yield. This result is consistent with the fact that the early bloom canopy spray of urea increased yield despite the fact that urea is not taken up by mature avocado leaves on the trees at the time of treatment.

In deciding whether to apply urea or boron during avocado bloom, the boron status of the orchard must be considered in order to avoid increasing leaf boron content to a toxic level. The results obtained in South Africa suggested that boron uptake by mature avocado leaves was limited. Our results demonstrated an average 10 ppm increase in leaf boron content in response to the early bloom boron spray.

If and when the application of either nutrient is made, it is critical to check and, if necessary, to adjust the pH of the spray solution to a value between pH 5.5 to 6.5 prior to spraying the trees to avoid damaging the flowers.

The author thanks the members of Dr. Ohr's and her laboratory for assisting with the harvests. Use of trade-names in this publication does not imply endorsement of the products named nor criticism of similar products not mentioned.

PARTNERS IN SUCCESS



and



- ▲ Worldwide marketing & sales
- ▲ Competitive grower returns
- ▲ Expert management, farming, & packing
- ▲ High per acre crop tonnages
- ▲ Effective and timely communication

**All this and more assures
you added profits. Let us show you how.**

CALL TODAY • 1-800-899-FARM • FAX 909-695-7593

SIERRA PACIFIC FARMS INC.

37 Years of Quality Service

BROKAW NURSERY, INC.



Avocado on Clonal and Seedling Rootstocks
Citrus • Kiwi • Cherimoya • Subtropicals
P.O. Box 4818, Saticoy, CA. 93004 • (805) 647-2262

If you buy a Spectrum...

**we'll earn money NOW,
you'll save money FROM NOW ON.**



**Citrus growers
can save
\$70-78/acre/yr
by cutting
chemicals • fuel • time.**

For information: Call Donald Cochran (805) 849-2624 or 849-3403 (after hours)

For more information circle #40 on reader service card

For more information circle #2 on reader service card

For more information circle #107 on reader service card

GROWER

CALIFORNIA

VOLUME 18, NO. 3 • MARCH 1994

f e a t u r e s

Still Growing 26

Hass Avocados Dominate Industry
by Howard Seelye

In The Beginning... 28

The Mother (Tree) Of All Hasses
by Howard Seelye

What Goes Around Comes Around 29

ACDI Builds Markets For U.S. Agriculture
by Bill Black

Healing The Healer 31

California's Workers' Comp Act Gets A Shot In The Arm
by Bob Kean

French Connection 37

A Look At The World's Leading Exporter
by James H. Hughes

regular columns

Agritutorial by Willard Thompson.....4
 An Apple A Day by Kenton Kidd7
 Perspectives by Mark Affleck12
 Mutual Concerns by Joel Nelsen19
 Watermark by Willard Thompson36

departments

Forum/Newsline5
 Apples7
 Trees & Vines10
 Avocados12
 Citrus19
 What's New33

This month's cover: As the avocado harvest moves into its peak season, CG looks at the variety that's making it all happen, starting on page 26. Photo by Willard Thompson.